

CSIR November 2020

Application No.	
Candidate Name	
Roll No.	
Test Date	26/11/2020
Test Time	3:00 PM - 6:00 PM
Subject	Mathematical Sciences

Section : **Part A Mathematical Sciences**

Q.1 Clock A loses 4 minutes every hour, clock B always shows the correct time and clock C gains 3 minutes every hour. On a Monday, all the three clocks showed the same time, 8 pm. On the following Wednesday, when the clock C shows 2 pm, what time will clock A show?

Options

1. 7: 20 am
2. 8: 40 am
3. 9:20 am
4. 10:40 am

Question Type : **MCQ**Question ID : **802437502**Option 1 ID : **8024372005**Option 2 ID : **8024372006**Option 3 ID : **8024372007**Option 4 ID : **8024372008**Status : **Not Answered**

Chosen Option : --

Q.2 A and B complete a work in 30 days. B and C complete the same work in 24 days whereas C and A complete the same work in 28 days. Based on these statements which of the following conclusions is correct?

Options 1.

C is the most efficient and B is the least efficient.

2.

B is the most efficient but, the least efficient one cannot be determined.

3.

C is the most efficient but, the least efficient one cannot be determined.

4.

C is the most efficient and A is the least efficient.

Question Type : **MCQ**

Question ID : **802437501**

Option 1 ID : **8024372001**

Option 2 ID : **8024372002**


Option 3 ID : **8024372003**

Option 4 ID : **8024372004**

Status : **Not Answered**

Chosen Option : **--**

- Q.3 The wavelength dependant absorbance of two compounds, A and B, is shown. Absorbance of mixture is a linear function of the concentration of the two compounds. R is defined as a ratio of absorbance at 650 nm to the absorbance at 950 nm.



If the mixture contains 95% of compound A then R must be

- Options
1. 95
 2. 5
 3. 1
 4. less than 1

Question Type : MCQ

Question ID : 802437499

Option 1 ID : 8024371993

Option 2 ID : 8024371994

Option 3 ID : 8024371995

Option 4 ID : 8024371996

Status : Not Answered

Chosen Option : --

Q.4

Find the value of $f(0)$ if $f(x+2) = (x+1)^{34} - (x+1)^{33} + 5$.

- Options
1. 5
 2. 7
 3. 6
 4. 72

Question Type : MCQ

Question ID : 802437491

Option 1 ID : 8024371961

Option 2 ID : 8024371962

Option 3 ID : 8024371963





Option 4 ID : 8024371964

Status : Answered

Chosen Option : 2

Q.5 An epidemic is spreading in a population of size P . The rate of spread R of the disease at a given time is proportional to both, the number of people affected by the disease (N), and the number of people not yet affected by the disease. Which of the following graphs of R vs N is correct?


Options



Question Type : MCQ
Question ID : 802437500
Option 1 ID : 8024371997

Option 2 ID : 8024371998
Option 3 ID : 8024371999
Option 4 ID : 8024372000
Status : Answered
Chosen Option : 4

Q.6 The shortest distance between the parallel lines A and B in the following figure is




Options

1. $\sqrt{2}$
2. 2
3. $2\sqrt{2}$
4. $2\sqrt{3}$

Question Type : MCQ
Question ID : 802437492
Option 1 ID : 8024371965
Option 2 ID : 8024371966
Option 3 ID : 8024371967
Option 4 ID : 8024371968
Status : Answered
Chosen Option : 1

- Q.7 A, B, C, D and E are the vertices of a regular pentagon as shown in the figure.



The angle $\angle ABC$ is

Options

1. 48°
2. 72°
3. 54°
4. 36°

Question Type : MCQ

Question ID : 802437509

Option 1 ID : 8024372033

Option 2 ID : 8024372034

Option 3 ID : 8024372035

Option 4 ID : 8024372036

Status : Not Answered

Chosen Option : --

- Q.8 A boat weighs 60 kg, and oarsmen A and B weigh 80 and 90 kg, respectively. Rowing at a constant power, the time required to complete a course is proportional to the total weight. Rowing alone, A and B complete the course in 1 and $1\frac{1}{2}$ hours, respectively. Assuming that their powers add up, how long will they take to complete the course if they row together?

Options

1. 49.4 min
2. 57.5 min
3. 62.6 min
4. 72.5 min

Question Type : MCQ

Question ID : 802437506

Option 1 ID : 8024372021

Option 2 ID : 8024372022


Option 3 ID : 8024372023

Option 4 ID : 8024372024

Status : Not Answered

Chosen Option : --

- Q.9 Consider a parallelogram ABCD with centre O and E as the midpoint of side CD. The area of the triangle OAE, is



Options

1. $\frac{1}{5}ah$
2. $\frac{1}{6}ah$
3. $\frac{1}{8}ah$
4. $\frac{1}{7}ah$

Question Type : MCQ

Question ID : 802437507

Option 1 ID : 8024372025

Option 2 ID : 8024372026

Option 3 ID : 8024372027

Option 4 ID : 8024372028


Status : Not Answered

Chosen Option : --


Q.10 Which is an appropriate diagram to represent the relations between the following categories: quadruped, mammal, whale, house lizard?

Options


1.




2.



3.



4.



Question Type : **MCQ**

Question ID : **802437495**

Option 1 ID : **8024371977**

Option 2 ID : **8024371978**

Option 3 ID : **8024371979**

Option 4 ID : **8024371980**

Status : **Not Answered**

Chosen Option : **--**

Q.11 Water is being filled in a cone from the top at a constant volumetric rate. The rate of increase of the height of the water column

Options

1. is linearly dependent on time.
2. depends on the apex angle of the cone.
3. increases as cube-root of the volumetric rate.
4. increases as square-root of the volumetric rate.

Question Type : **MCQ**

Question ID : **802437497**

Option 1 ID : **8024371985**

Option 2 ID : **8024371986**

Option 3 ID : **8024371987**

Option 4 ID : **8024371988**

Status : **Not Answered**

Chosen Option : --

Q.12 The given table shows the numbers of active and recovered cases of a certain disease. Assuming that the linear trend for both continues, on which day will recovered cases be twice that of the active cases?

Day	0	1	4	7	10
Active cases	990	1000	1030	1060	1090
Recovered cases	760	800	920	1040	1160

Options

1. 61
2. 62
3. 63
4. 64

Question Type : **MCQ**

Question ID : **802437505**

Option 1 ID : **8024372017**

Option 2 ID : **8024372018**

Option 3 ID : **8024372019**

Option 4 ID : **8024372020**

Status : **Not Answered**

Chosen Option : --

Q.13 Two varieties A and B of rice cost Rs. 30 and Rs. 90 per kg, whereas two varieties C and D of pulses, Rs. 100 and Rs. 120 per kg, respectively. If at least one kg each of A and B and at least half a kg each of C and D have to be purchased, then the minimum and maximum costs of a total of 5 kg of these provisions are, respectively

Options

1. Rs. 150 and Rs. 600
2. Rs. 260 and Rs. 530
3. Rs. 290 and Rs. 470
4. Rs. 370 and Rs. 460

Question Type : **MCQ**

Question ID : **802437493**

Option 1 ID : **8024371969**

Option 2 ID : **8024371970**

Option 3 ID : **8024371971**

Option 4 ID : **8024371972**

Status : **Not Answered**

Chosen Option : --

Q.14 One of four suspects A, B, C and D has committed a crime. A and D are always truthful, and B and C are always untruthful. C and D are identical twins and the interrogator does not know who is who. If A says, "D is innocent", B says, "A is guilty" and among C and D one says, "A is innocent" and the other says, "B is guilty", then which of the following is FALSE?

Options

1. D said "A is innocent"
2. D is innocent
3. B is innocent
4. C is innocent

Question Type : **MCQ**

Question ID : **802437494**

Option 1 ID : **8024371973**

Option 2 ID : **8024371974**

Option 3 ID : **8024371975**

Option 4 ID : **8024371976**

Status : **Not Answered**

Chosen Option : --

Q.15 A 7 m long tube having inner diameter of 2 cm is filled with water. The water is then poured into a cylindrical bucket having inner base area of 200 cm^2 . What will be the approximate height (in cm) of water in the bucket?

Options

1. 22
2. 44
3. 9
4. 11

Question Type : **MCQ**

Question ID : **802437496**

Option 1 ID : **8024371981**

Option 2 ID : **8024371982**

Option 3 ID : **8024371983**

Option 4 ID : **8024371984**

Status : **Answered**

Chosen Option : **2**

Q.16 On a 200 m long straight road, maximum number of poles are fixed at 20 m interval. How many of these poles should be removed in order to have maximum number of poles at an interval of 40 m on the road?

Options

1. 8
2. 6
3. 5
4. 4

Question Type : **MCQ**

Question ID : **802437510**

Option 1 ID : **8024372037**

Option 2 ID : **8024372038**

Option 3 ID : **8024372039**

Option 4 ID : **8024372040**

Status : **Answered**

Chosen Option : **3**

Q.17 In a class, there is one pencil for every two students, one eraser for every three students, and one ruler for every four students. If the total number of these stationery items required is 65, how many students are present in the class?

Options

1. 55
2. 60
3. 65
4. 70

Question Type : **MCQ**

Question ID : **802437503**

Option 1 ID : **8024372009**

Option 2 ID : **8024372010**

Option 3 ID : **8024372011**

Option 4 ID : **8024372012**

Status : **Not Answered**

Chosen Option : --

Q.18 A square board is divided into 9 smaller identical squares by drawing lines. Three bullets are shot at the board randomly. The probability that at least 2 bullets hit the same small square is,

Options

1. $1/3$
2. $56/81$
3. $25/81$
4. $2/3$

Question Type : **MCQ**

Question ID : **802437498**

Option 1 ID : **8024371989**

Option 2 ID : **8024371990**


Option 3 ID : **8024371991**

Option 4 ID : **8024371992**

Status : **Not Answered**

Chosen Option : --

- Q.19 The figure shows temperature and salinity of four samples of water.
Which one of the samples has the highest density?



Options

1. A
2. B
3. C
4. D

Question Type : MCQ

Question ID : 802437504

Option 1 ID : 8024372013

Option 2 ID : 8024372014

Option 3 ID : 8024372015

Option 4 ID : 8024372016

Status : Answered

Chosen Option : 1

- Q.20 The sum of the first n even numbers is

Options

1. divisible by n and not by $(n + 1)$
2. divisible by $(n + 1)$ and not by n
3. divisible by both n and $(n + 1)$
4. neither divisible by n nor by $(n + 1)$

Question Type : MCQ

Question ID : 802437508

Option 1 ID : 8024372029

Option 2 ID : 8024372030

Option 3 ID : 8024372031

Option 4 ID : 8024372032

Status : Answered

Chosen Option : 3

Q.1 Let A be a 2×2 real matrix with $\det A = 1$ and $\text{trace } A = 3$. What is the value of $\text{trace } A^2$?

Options

1. 2
2. 10
3. 9
4. 7

Question Type : **MCQ**

Question ID : **802437520**

Option 1 ID : **8024372077**

Option 2 ID : **8024372078**

Option 3 ID : **8024372079**

Option 4 ID : **8024372080**

Status : **Answered**

Chosen Option : **4**

Q.2 Given f, g are continuous functions on $[0,1]$ such that $f(0) = f(1) = 0$; $g(0) = g(1) = 1$ and $f(1/2) > g(1/2)$. Which of the following statements is true?

Options

1. There is no $t \in [0, 1]$ such that $f(t) = g(t)$
2. There is exactly one $t \in [0, 1]$ such that $f(t) = g(t)$
3. There are at least two $t \in [0, 1]$ such that $f(t) = g(t)$
4. There are always infinitely many $t \in [0, 1]$ such that $f(t) = g(t)$

Question Type : **MCQ**

Question ID : **802437516**

Option 1 ID : **8024372061**

Option 2 ID : **8024372062**

Option 3 ID : **8024372063**

Option 4 ID : **8024372064**

Status : **Answered**

Chosen Option : **4**

Q.3

Which of the following statements is true?

Options 1.

1. There are at most countably many continuous maps from \mathbb{R}^2 to \mathbb{R} .
2. There are at most finitely many continuous surjective maps from \mathbb{R}^2 to \mathbb{R} .
3. There are infinitely many continuous injective maps from \mathbb{R}^2 to \mathbb{R} .
4. There are no continuous bijective maps from \mathbb{R}^2 to \mathbb{R} .

Question Type : MCQ

Question ID : 802437513

Option 1 ID : 8024372049

Option 2 ID : 8024372050

Option 3 ID : 8024372051

Option 4 ID : 8024372052

Status : Answered

Chosen Option : 4

Q.4

Let A and B be 2×2 matrices. Then which of the following is true?

Options

1. $\det(A + B) + \det(A - B) = \det A + \det B$
2. $\det(A + B) + \det(A - B) = 2\det A - 2\det B$
3. $\det(A + B) + \det(A - B) = 2\det A + 2\det B$
4. $\det(A + B) - \det(A - B) = 2\det A - 2\det B$

Question Type : MCQ

Question ID : 802437518

Option 1 ID : 8024372069

Option 2 ID : 8024372070

Option 3 ID : 8024372071

Option 4 ID : 8024372072

Status : Answered

Chosen Option : 3

Q.5

The series

$$\sum_{n=1}^{\infty} \frac{(-1)^n \sin nx}{n^{\log_e n}}, \quad x \in \mathbb{R}$$

converges

Options

1. only for $x = 0$
2. uniformly only for $x \in [-\pi, \pi]$
3. uniformly only for $x \in \mathbb{R} \setminus \{n\pi : n \in \mathbb{Z}\}$
4. uniformly for all $x \in \mathbb{R}$

Question Type : **MCQ**Question ID : **802437514**Option 1 ID : **8024372053**Option 2 ID : **8024372054**Option 3 ID : **8024372055**Option 4 ID : **8024372056**Status : **Not Answered**

Chosen Option : --

Q.6

Let $\{E_n\}$ be a sequence of subsets of \mathbb{R} .

Define

$$\limsup_n E_n = \bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} E_n$$

$$\liminf_n E_n = \bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} E_n$$

Which of the following statements is true?

Options

1. $\limsup_n E_n = \liminf_n E_n$
2. $\limsup_n E_n = \{x: x \in E_n \text{ for some } n\}$
3. $\liminf_n E_n = \{x: x \in E_n \text{ for all but finitely many } n\}$
4. $\liminf_n E_n = \{x: x \in E_n \text{ for infinitely many } n\}$

Question Type : MCQ

Question ID : 802437511

Option 1 ID : 8024372041

Option 2 ID : 8024372042

Option 3 ID : 8024372043

Option 4 ID : 8024372044

Status : Answered

Chosen Option : 1

Q.7 Which of the following real quadratic forms on \mathbb{R}^2 is positive definite?

Options

1. $Q(X, Y) = XY$
2. $Q(X, Y) = X^2 - XY + Y^2$
3. $Q(X, Y) = X^2 + 2XY + Y^2$
4. $Q(X, Y) = X^2 + XY$

Question Type : **MCQ**

Question ID : **802437522**

Option 1 ID : **8024372085**

Option 2 ID : **8024372086**

Option 3 ID : **8024372087**

Option 4 ID : **8024372088**

Status : **Answered**

Chosen Option : **2**

Q.8 Let A be an $n \times n$ matrix such that the set of all its nonzero eigenvalues has exactly r elements. Which of the following statements is true?

Options

1. $\text{rank } A \leq r$
2. If $r = 0$, then $\text{rank } A < n - 1$
3. $\text{rank } A \geq r$
4. A^2 has r distinct nonzero eigenvalues

Question Type : **MCQ**

Question ID : **802437517**

Option 1 ID : **8024372065**

Option 2 ID : **8024372066**

Option 3 ID : **8024372067**

Option 4 ID : **8024372068**

Status : **Answered**

Chosen Option : **2**

Q.9 $f: \mathbb{N} \rightarrow \mathbb{N}$ be a bounded function. Which of the following statements is NOT true?

Options

1. $\limsup_{n \rightarrow \infty} f(n) \in \mathbb{N}$
2. $\liminf_{n \rightarrow \infty} f(n) \in \mathbb{N}$
3. $\liminf_{n \rightarrow \infty} (f(n) + n) \in \mathbb{N}$
4. $\limsup_{n \rightarrow \infty} (f(n) + n) \notin \mathbb{N}$

Question Type : **MCQ**

Question ID : **802437512**

Option 1 ID : **8024372045**

Option 2 ID : **8024372046**

Option 3 ID : **8024372047**

Option 4 ID : **8024372048**

Status : **Answered**

Chosen Option : **3**

Q.10

If $A = \begin{pmatrix} 3 & -2 \\ 2 & -1 \end{pmatrix}$, then A^{20} equals

Options

1. $\begin{pmatrix} 41 & 40 \\ -40 & -39 \end{pmatrix}$
2. $\begin{pmatrix} 41 & -40 \\ 40 & -39 \end{pmatrix}$
3. $\begin{pmatrix} 41 & -40 \\ -40 & -39 \end{pmatrix}$
4. $\begin{pmatrix} 41 & 40 \\ 40 & -39 \end{pmatrix}$

Question Type : **MCQ**

Question ID : **802437519**

Option 1 ID : **8024372073**

Option 2 ID : **8024372074**

Option 3 ID : **8024372075**

Option 4 ID : **8024372076**

Status : **Answered**

Chosen Option : **2**

Q.11 For $a, b \in \mathbb{R}$, let

$$p(x, y) = a^2x_1y_1 + abx_2y_1 + abx_1y_2 + b^2x_2y_2, \quad x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{R}^2.$$

For what values of a and b does $p: \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ define an inner product?

Options

1. $a > 0, b > 0$
2. $ab > 0$
3. $a = 0, b = 0$
4. For no values of a, b

Question Type : MCQ

Question ID : 802437521

Option 1 ID : 8024372081

Option 2 ID : 8024372082

Option 3 ID : 8024372083

Option 4 ID : 8024372084

Status : Answered

Chosen Option : 2

Q.12 Given $(a_n)_{n \geq 1}$ a sequence of real numbers, which of the following statements is true?

Options

1. $\sum_{n \geq 1} (-1)^n \frac{a_n}{1 + |a_n|}$ converges

2.

There is a subsequence $(a_{n_k})_{k \geq 1}$ such that $\sum_{k \geq 1} \frac{a_{n_k}}{1 + |a_{n_k}|}$ converges

3.

There is a number b such that $\sum_{n \geq 1} \left| b - \frac{a_n}{1 + |a_n|} \right| (-1)^n$ converges

4.

There is a number b and a subsequence $(a_{n_k})_{k \geq 1}$ such that

$$\sum_{k \geq 1} \left| b - \frac{a_{n_k}}{1 + |a_{n_k}|} \right| \text{ converges}$$

Question Type : MCQ

Question ID : 802437515

Option 1 ID : 8024372057

Option 2 ID : 8024372058

Option 3 ID : 8024372059

Option 4 ID : 8024372060

Status : Answered

Chosen Option : 1

Q.13

Let $\varphi(n)$ be the cardinality of the set

$$\{a \mid 1 \leq a \leq n, (a, n) = 1\}$$

where (a, n) denotes the *gcd* of a and n . Which of the following is NOT true?

Options 1.

1. There exist infinitely many n such that $\varphi(n) > \varphi(n + 1)$.

2.

2. There exist infinitely many n such that $\varphi(n) < \varphi(n + 1)$.

3.

3. There exists $N \in \mathbb{N}$ such that $N > 2$ and for all $n > N$,

$$\varphi(N) < \varphi(n)$$

4.

4. The set $\left\{\frac{\varphi(n)}{n} : n \in \mathbb{N}\right\}$ has finitely many limit points.

Question Type : MCQ

Question ID : 802437529

Option 1 ID : 8024372113

Option 2 ID : 8024372114

Option 3 ID : 8024372115

Option 4 ID : 8024372116

Status : Answered

Chosen Option : 3

Q.14

For a positive integer p , consider the holomorphic function

$$f(z) = \frac{\sin z}{z^p} \text{ for } z \in \mathbb{C} \setminus \{0\}.$$

For which values of p does there exist a holomorphic function $g : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$ such that $f(z) = g'(z)$ for $z \in \mathbb{C} \setminus \{0\}$?

Options

1. All even integers

2. All odd integers

3. All multiples of 3

4. All multiples of 4

Question Type : MCQ

Question ID : 802437524

Option 1 ID : 8024372093

Option 2 ID : 8024372094

Option 3 ID : 8024372095

Option 4 ID : 8024372096

Status : Not Answered

Chosen Option : --

- Q.15** Let p be a positive integer. Consider the closed curve $r(t) = e^{it}, 0 \leq t < 2\pi$. Let f be a function holomorphic in $\{z: |z| < R\}$ where $R > 1$. If f has a zero only at $z_0, 0 < |z_0| < R$, and it is of multiplicity q , then

$$\frac{1}{2\pi i} \int_r \frac{f'(z)}{f(z)} z^p dz \text{ equals}$$

Options

1. qz_0^p
2. z_0q^p
3. pz_0^q
4. z_0p^q

Question Type : **MCQ**

Question ID : **802437526**

Option 1 ID : **8024372101**

Option 2 ID : **8024372102**

Option 3 ID : **8024372103**

Option 4 ID : **8024372104**

Status : **Answered**

Chosen Option : 1

- Q.16** For any two metric spaces $(X, d_X), (Y, d_Y)$ a map $f: X \rightarrow Y$ is said to be a closed map if whenever F is closed in X , then $f(F)$ is closed in Y . For any subset B of a metric space, B is given the induced metric. The metric on $X \times Y$ is given by $d((x, y), (x', y')) = \max\{d_X(x, x'), d_Y(y, y')\}$. Which of the following are true?

Options 1.

For any subset $A \subseteq X$ the inclusion map $i: A \rightarrow X$ is closed

2.

The projection map $p_1: X \times Y \rightarrow X$ given by $p_1(x, y) = x$ is closed

3.

Suppose that $f: X \rightarrow Y, g: Y \rightarrow Z$ are continuous maps. If $g \circ f: X \rightarrow Z$ is a closed map then $g|_{f(X)}: f(X) \rightarrow Z$ is closed. Here $g|_{f(X)}$ means the map g restricted to $f(X)$

4.

If $f: X \rightarrow Y$ takes closed balls into closed sets then f is closed

Question Type : **MCQ**

Question ID : **802437530**

Option 1 ID : **8024372117**

Option 2 ID : **8024372118**

Option 3 ID : **8024372119**

Option 4 ID : **8024372120**

Status : **Not Answered**

Chosen Option : --

Q.17 Let X be a non-empty set and $P(X)$ be the set of all subsets of X . On $P(X)$, define two operations $*$ and Δ as follows: for $A, B \in P(X)$, $A * B = A \cap B$; $A \Delta B = (A \cup B) \setminus (A \cap B)$.

Which of the following statements is true?

Options

1. $P(X)$ is a group under $*$ as well as under Δ
2. $P(X)$ is a group under $*$, but not under Δ
3. $P(X)$ is a group under Δ , but not under $*$
4. $P(X)$ is neither a group under $*$ nor under Δ

Question Type : **MCQ**

Question ID : **802437528**

Option 1 ID : **8024372109**

Option 2 ID : **8024372110**

Option 3 ID : **8024372111**

Option 4 ID : **8024372112**

Status : **Answered**

Chosen Option : **3**

Q.18

Which of the following statements is true?

Options

1. Every even integer $n \geq 16$ divides $(n - 1)! + 3$
2. Every odd integer $n \geq 16$ divides $(n - 1)!$
3. Every even integer $n \geq 16$ divides $(n - 1)!$
4. For every integer $n \geq 16$, n^2 divides $n! + 1$

Question Type : **MCQ**

Question ID : **802437527**

Option 1 ID : **8024372105**

Option 2 ID : **8024372106**

Option 3 ID : **8024372107**

Option 4 ID : **8024372108**

Status : **Answered**

Chosen Option : **3**

Q.19 Let γ be the positively oriented circle in the complex plane given by $\{z \in \mathbb{C} : |z - 1| = 1\}$. Then

$$\frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z^3 - 1} \text{ equals}$$

Options

1. 3
2. 1/3
3. 2
4. 1/2

Question Type : MCQ

Question ID : 802437523

Option 1 ID : 8024372089

Option 2 ID : 8024372090

Option 3 ID : 8024372091

Option 4 ID : 8024372092

Status : Answered

Chosen Option : 2

Q.20 Let γ be the positively oriented circle in the complex plane given by $\{z \in \mathbb{C} : |z - 1| = 1/2\}$. The line integral

$$\int_{\gamma} \frac{ze^{1/z}}{z^2 - 1} dz$$

equals

Options

1. $i\pi e$
2. $-i\pi e$
3. πe
4. $-\pi e$

Question Type : MCQ

Question ID : 802437525

Option 1 ID : 8024372097

Option 2 ID : 8024372098

Option 3 ID : 8024372099

Option 4 ID : 8024372100

Status : Answered

Chosen Option : 1

Q.21 The general solution of the surfaces which are perpendicular to the family of surfaces

$$z^2 = kxy, k \in \mathbb{R}$$

is

Options

1. $\phi(x^2 - y^2, xz) = 0, \phi \in C^1(\mathbb{R}^2)$
2. $\phi(x^2 - y^2, x^2 + z^2) = 0, \phi \in C^1(\mathbb{R}^2)$
3. $\phi(x^2 - y^2, 2x^2 + z^2) = 0, \phi \in C^1(\mathbb{R}^2)$
4. $\phi(x^2 + y^2, 3x^2 - z^2) = 0, \phi \in C^1(\mathbb{R}^2)$

Question Type : **MCQ**

Question ID : **802437533**

Option 1 ID : **8024372129**

Option 2 ID : **8024372130**

Option 3 ID : **8024372131**

Option 4 ID : **8024372132**

Status : **Not Answered**

Chosen Option : --

Q.22 Let f be an infinitely differentiable real-valued function on a bounded interval I . Take $n \geq 1$ interpolation points $\{x_0, x_1, \dots, x_{n-1}\}$. Take n additional interpolation points

$$x_{n+j} = x_j + \varepsilon, \quad j = 0, 1, \dots, n-1$$

where $\varepsilon > 0$ is such that $\{x_0, x_1, \dots, x_{2n-1}\}$ are all distinct.

Let p_{2n-1} be the Lagrange interpolation polynomial of degree $2n-1$ with the interpolation points $\{x_0, x_1, \dots, x_{2n-1}\}$ for the function f .

Let q_{2n-1} be the Hermite interpolation polynomial of degree $2n-1$ with the interpolation points $\{x_0, x_1, \dots, x_{n-1}\}$ for the function f . In the $\varepsilon \rightarrow 0$ limit, the quantity

$$\sup_{x \in I} |p_{2n-1}(x) - q_{2n-1}(x)|$$

Options

1. does not necessarily converge
2. converges to $\frac{1}{2n}$
3. converges to 0
4. converges to $\frac{1}{2n+1}$

Question Type : **MCQ**

Question ID : **802437535**

Option 1 ID : **8024372137**

Option 2 ID : **8024372138**

Option 3 ID : **8024372139**

Option 4 ID : **8024372140**

Status : **Not Answered**

Chosen Option : --

Q.23 Let $y_0 > 0, z_0 > 0$ and $\alpha > 1$.

Consider the following two differential equations:

$$(*) \begin{cases} \frac{dy}{dt} = y^\alpha & \text{for } t > 0, \\ y(0) = y_0 \end{cases}$$

$$(**) \begin{cases} \frac{dz}{dt} = -z^\alpha & \text{for } t > 0, \\ z(0) = z_0 \end{cases}$$

We say that the solution to a differential equation exists globally if it exists for all $t > 0$.

Which of the following statements is true?

Options

1. Both (*) and (**) have global solutions
2. None of (*) and (**) have global solutions

3.

There exists a global solution for (*) and there exists a $T < \infty$ such that

$$\lim_{t \rightarrow T} |z(t)| = +\infty$$

4.

There exists a global solution for (**) and there exists a $T < \infty$ such that

$$\lim_{t \rightarrow T} |y(t)| = +\infty$$

Question Type : **MCQ**

Question ID : **802437532**

Option 1 ID : **8024372125**

Option 2 ID : **8024372126**

Option 3 ID : **8024372127**

Option 4 ID : **8024372128**

Status : **Not Answered**

Chosen Option : --

Q.24

The solution of the Fredholm integral equation

$$y(s) = s + 2 \int_0^1 (st^2 + s^2t)y(t)dt$$

is

Options

1. $y(s) = -(50s + 40s^2)$
2. $y(s) = (30s + 15s^2)$
3. $y(s) = -(30s + 40s^2)$
4. $y(s) = (60s + 50s^2)$

Question Type : MCQ

Question ID : 802437537

Option 1 ID : 8024372145

Option 2 ID : 8024372146

Option 3 ID : 8024372147

Option 4 ID : 8024372148

Status : Not Attempted and
Marked For Review

Chosen Option : --

Q.25

Let k be a positive integer. Consider the differential equation

$$\begin{cases} \frac{dy}{dt} = y^{\frac{5k}{5k+2}} \text{ for } t > 0, \\ y(0) = 0 \end{cases}$$

Which of the following statements is true?

Options 1.

1. It has a unique solution which is continuously differentiable on $(0, \infty)$
2. It has at most two solutions which are continuously differentiable on $(0, \infty)$
3. It has infinitely many solutions which are continuously differentiable on $(0, \infty)$
4. It has no continuously differentiable solution on $(0, \infty)$

Question Type : MCQ

Question ID : 802437531

Option 1 ID : 8024372121

Option 2 ID : 8024372122

Option 3 ID : 8024372123

Option 4 ID : 8024372124

Status : Not Answered

Chosen Option : --

Q.26 The extremal of the functional

$$J(y) = \int_0^1 [2(y')^2 + xy]dx, \quad y(0) = 0, y(1) = 1, y \in C^2[0, 1]$$

is

Options

1. $y = \frac{x^2}{12} + \frac{11x}{12}$

2. $y = \frac{x^3}{3} + \frac{2x^2}{3}$

3. $y = \frac{x^2}{7} + \frac{6x}{7}$

4. $y = \frac{x^3}{24} + \frac{23x}{24}$

Question Type : MCQ

Question ID : 802437536

Option 1 ID : 8024372141

Option 2 ID : 8024372142

Option 3 ID : 8024372143

Option 4 ID : 8024372144

Status : Answered

Chosen Option : 4

Q.27 The general solution of the equation

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 0$$

is

Options

1. $z = \phi\left(\frac{|x|}{|y|}\right), \phi \in C^1(\mathbb{R})$

2. $z = \phi\left(\frac{x-1}{y}\right), \phi \in C^1(\mathbb{R})$

3. $z = \phi\left(\frac{x+1}{y}\right), \phi \in C^1(\mathbb{R})$

4. $z = \phi(|x| + |y|), \phi \in C^1(\mathbb{R})$

Question Type : MCQ

Question ID : 802437534

Option 1 ID : 8024372133

Option 2 ID : 8024372134

Option 3 ID : 8024372135

Option 4 ID : 8024372136

Status : Answered

Chosen Option : 1

Q.28 Consider the solid S made of a material of constant density in the shape of a hemisphere of unit radius:

$$S = \{(x, y, z) | x^2 + y^2 + z^2 \leq 1 \quad z \geq 0\}.$$

Which of the following statements is true?

Options

1. The centre of mass of S is at the origin
2. The x -axis is a principal axis for S
3. The moment of inertia tensor of S is not a diagonal matrix
4. The z -axis is a principal axis for S

Question Type : **MCQ**

Question ID : **802437538**

Option 1 ID : **8024372149**

Option 2 ID : **8024372150**

Option 3 ID : **8024372151**

Option 4 ID : **8024372152**

Status : **Answered**

Chosen Option : 1

Q.29 The maximum and the minimum values of $5x + 7y$, when $|x| + |y| \leq 1$ are

Options

1. 5 and -5
2. 5 and -7
3. 7 and -5
4. 7 and -7

Question Type : **MCQ**

Question ID : **802437550**

Option 1 ID : **8024372197**

Option 2 ID : **8024372198**

Option 3 ID : **8024372199**

Option 4 ID : **8024372200**

Status : **Answered**

Chosen Option : 4

Q.30 Consider a Markov Chain X_0, X_1, X_2, \dots with state space S . Suppose $i, j \in S$ are two states which communicate with each other. Which of the following statements is NOT correct?

Options

1. Period of i = period of j
2. i is recurrent if and only if j is recurrent
3. $\lim_{n \rightarrow \infty} P(X_n = i | X_0 = k) = \lim_{n \rightarrow \infty} P(X_n = j | X_0 = k)$ for all $k \in S$
4. $\lim_{n \rightarrow \infty} P(X_n = j | X_0 = i) = \lim_{n \rightarrow \infty} P(X_n = j | X_0 = j)$

Question Type : **MCQ**

Question ID : **802437541**

Option 1 ID : **8024372161**

Option 2 ID : **8024372162**

Option 3 ID : **8024372163**

Option 4 ID : **8024372164**

Status : **Answered**

Chosen Option : **3**

Q.31 Let X_1 and X_2 be two i.i.d. $p \times 1$ multivariate normal random vectors with mean μ and positive definite dispersion matrix Σ . Then which of the following random variables always has a central chi-square distribution

Options

1. $\frac{1}{2}(X_1 - X_2)^T(X_1 - X_2)$
2. $2(X_1 - X_2)^T(X_1 - X_2)$
3. $2(X_1 - X_2)^T \Sigma^{-1}(X_1 - X_2)$
4. $\frac{1}{2}(X_1 - X_2)^T \Sigma^{-1}(X_1 - X_2)$

Question Type : **MCQ**

Question ID : **802437547**

Option 1 ID : **8024372185**

Option 2 ID : **8024372186**

Option 3 ID : **8024372187**

Option 4 ID : **8024372188**

Status : **Not Answered**

Chosen Option : **--**

Q.32 Suppose that X has uniform distribution on the interval $[0, 100]$. Let Y denote the greatest integer smaller than or equal to X . Which of the following is true?

Options

1. $P(Y \leq 25) = 1/4$
2. $P(Y \leq 25) = 26/100$
3. $E(Y) = 50$
4. $E(Y) = 101/2$

Question Type : **MCQ**

Question ID : **802437542**

Option 1 ID : **8024372165**

Option 2 ID : **8024372166**

Option 3 ID : **8024372167**

Option 4 ID : **8024372168**

Status : **Not Answered**

Chosen Option : --

Q.33 10 units are chosen by simple random sampling without replacement from a population of size 100. Consider the sample variance $\frac{1}{10} \sum_{i=1}^{10} (y_i - \bar{y})^2 = s^2$. An unbiased estimate of population variance $\sigma^2 = \frac{1}{100} \sum_{i=1}^{100} (Y_i - \bar{Y})^2$ is

Options

1. s^2
2. $\frac{10}{11} s^2$
3. $\frac{100}{99} s^2$
4. $\frac{100}{111} s^2$

Question Type : **MCQ**

Question ID : **802437548**

Option 1 ID : **8024372189**

Option 2 ID : **8024372190**

Option 3 ID : **8024372191**

Option 4 ID : **8024372192**

Status : **Not Answered**

Chosen Option : --

Q.34 Consider a Randomized Block Design with b blocks and k treatments. Let the observation corresponding to the i^{th} treatment and the j^{th} block be y_{ij} , $1 \leq i \leq k$, $1 \leq j \leq b$, which satisfies the usual linear model. Which of the following is true?

Options 1.

1. The estimates of any two treatment contrasts are uncorrelated
2. The error sum of squares has $bk - 1$ degrees of freedom
3. The estimate of any treatment contrast is uncorrelated with the estimate of any block contrast
4. The correlation between the estimates of two treatment contrasts is always negative

Question Type : **MCQ**
Question ID : **802437549**
Option 1 ID : **8024372193**
Option 2 ID : **8024372194**
Option 3 ID : **8024372195**
Option 4 ID : **8024372196**
Status : **Not Answered**
Chosen Option : --

Q.35 Consider 35 i.i.d. observations X_1, X_2, \dots, X_{15} and Y_1, Y_2, \dots, Y_{20} . Let R be the Wilcoxon's rank sum statistic based on the ranks of the X 's in the combined sample. Then the expected value of R is

Options

1. 270
2. 300
3. 360.5
4. 330.5

Question Type : **MCQ**
Question ID : **802437545**
Option 1 ID : **8024372177**
Option 2 ID : **8024372178**
Option 3 ID : **8024372179**
Option 4 ID : **8024372180**
Status : **Not Answered**
Chosen Option : --

Q.36 Consider the pdf given by

$$f(x|\theta) = \frac{e^{-(x-\theta)}}{[1+e^{-(x-\theta)}]^2}, \quad -\infty < x < \infty, \quad -\infty < \theta < \infty$$

Based on one observation X with the above pdf, a UMP test of size α for testing

$H_0: \theta \leq \theta_0$ versus $H_1: \theta > \theta_0$ is

Options

1. $X > k$ for some k such that $\alpha = P_{\theta_0}[X > k]$
2. $X < k$ for some k such that $\alpha = P_{\theta_0}[X > k]$
3. $X > k$ for some k such that $\alpha = P_{\theta_0}[X < k]$
4. $X < k$ for some k such that $\alpha = P_{\theta_0}[X < k]$

Question Type : **MCQ**

Question ID : **802437544**

Option 1 ID : **8024372173**

Option 2 ID : **8024372174**

Option 3 ID : **8024372175**

Option 4 ID : **8024372176**

Status : **Not Answered**

Chosen Option : --

Q.37

In an examination involving multiple choice questions, a student works out the solution in 50% of the questions. In the remaining questions the student guesses the answer. However, when the answer is guessed the probability that it is correct is 0.30. When the student works out the solutions it may be wrong with probability 0.10.

If the answer to a particular question is correct, what is the probability that the student guessed the answer?

Options

1. **0.25**
2. **0.50**
3. **0.90**
4. **0.30**

Question Type : **MCQ**

Question ID : **802437539**

Option 1 ID : **8024372153**

Option 2 ID : **8024372154**

Option 3 ID : **8024372155**

Option 4 ID : **8024372156**

Status : **Not Answered**

Chosen Option : --

Q.38 Let X_1, X_2, \dots, X_n be i.i.d. random variables with common pdf
 $f(x|\theta) = \frac{(\log \theta)\theta^x}{\theta - 1}$, for
 $0 < x < 1$ where $\theta > 1$ is an unknown parameter. Then the statistic
 $T = \sum_{i=1}^n X_i$ is

- Options
1. sufficient, but not complete
 2. sufficient, but not minimal sufficient
 3. complete sufficient
 4. neither complete, nor sufficient

Question Type : **MCQ**
 Question ID : **802437543**
 Option 1 ID : **8024372169**
 Option 2 ID : **8024372170**
 Option 3 ID : **8024372171**
 Option 4 ID : **8024372172**
 Status : **Not Answered**
 Chosen Option : --

Q.39 Let X_1, X_2, \dots be i.i.d. random variables having a χ^2 -distribution with 5 degrees of freedom.

Let $a \in \mathbb{R}$ be constant. Then the limiting distribution of $a \left(\frac{X_1 + \dots + X_n - 5n}{\sqrt{n}} \right)$ is

- Options 1.
1. Gamma distribution for an appropriate value of a
 2. χ^2 -distribution for an appropriate value of a
 3. Standard normal distribution for an appropriate value of a
 4. A degenerate distribution for an appropriate value of a

Question Type : **MCQ**
 Question ID : **802437540**
 Option 1 ID : **8024372157**
 Option 2 ID : **8024372158**
 Option 3 ID : **8024372159**
 Option 4 ID : **8024372160**
 Status : **Not Answered**
 Chosen Option : --

Q.40 Let $I, J > 5$. Consider two-way ANOVA, where the observations satisfy the linear model

$$y_{ij} = \alpha + \beta_i + \gamma_j + \varepsilon_{ij}, \quad 1 \leq i \leq I, \quad 1 \leq j \leq J$$

$$E(\varepsilon_{ij}) = 0, \text{Var}(\varepsilon_{ij}) = \sigma^2, \quad \sum_{i=1}^I \beta_i = \sum_{j=1}^J \gamma_j = 0.$$

In this set-up

Options

1. β_1 is estimable
2. γ_1 is estimable
3. $\beta_1 - \beta_2$ is estimable
4. $\beta_1 + \gamma_2$ is estimable

Question Type : **MCQ**

Question ID : **802437546**

Option 1 ID : **8024372181**

Option 2 ID : **8024372182**

Option 3 ID : **8024372183**

Option 4 ID : **8024372184**

Status : **Not Answered**

Chosen Option : **--**

Section : **Part C Mathematical Sciences**

Q.1

Let $p: \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$p(x, y) = \begin{cases} |x| & \text{if } x \neq 0 \\ |y| & \text{if } x = 0. \end{cases}$$

Which of the following statements are true?

Options

1. $p(x, y) = 0$ if and only if $x = y = 0$
2. $p(x, y) \geq 0$ for all x, y
- 3.
4. $p(ax, ay) = |a| p(x, y)$ for all $a \in \mathbb{R}$ and for all x, y
4. $p(x_1 + x_2, y_1 + y_2) \leq p(x_1, y_1) + p(x_2, y_2)$ for all $(x_1, y_1), (x_2, y_2)$

Question Type : **MSQ**

Question ID : **802437560**

Option 1 ID : **8024372237**

Option 2 ID : **8024372238**

Option 3 ID : **8024372239**

Option 4 ID : **8024372240**

Status : **Answered**

Chosen Option : **1,2,3**

Q.2 Let $Q(x, y, z)$ be a real quadratic form. Which of the following statements are true?

Options 1.

1. $Q(x_1 + x_2, y, z) = Q(x_1, y, z) + Q(x_2, y, z)$ for all x_1, x_2, y, z

2.

$Q(x_1 + x_2, y_1 + y_2, 0) + Q(x_1 - x_2, y_1 - y_2, 0) = 2Q(x_1, y_1, 0) + 2Q(x_2, y_2, 0)$ for all x_1, x_2, y_1, y_2

3.

$Q(x_1 + x_2, y_1 + y_2, z_1 + z_2) = Q(x_1, y_1, z_1) + Q(x_2, y_2, z_2)$
for at least one choice of $x_1, x_2, y_1, y_2, z_1, z_2$

4.

$2Q(x_1 + x_2, y_1 + y_2, 0) + 2Q(x_1 - x_2, y_1 - y_2, 0) = Q(x_1, y_1, 0) + Q(x_2, y_2, 0)$ for all x_1, x_2, y_1, y_2

Question Type : **MSQ**

Question ID : **802437568**

Option 1 ID : **8024372269**

Option 2 ID : **8024372270**

Option 3 ID : **8024372271**

Option 4 ID : **8024372272**

Status : **Not Attempted and Marked For Review**

Chosen Option : --

Q.3 Let P be a square matrix such that $P^2 = P$. Which of the following statements are true?

Options

1. Trace of P is an irrational number
2. Trace of $P = \text{rank of } P$
3. Trace of P is an integer
4. Trace of P is an imaginary complex number

Question Type : **MSQ**

Question ID : **802437561**

Option 1 ID : **8024372241**

Option 2 ID : **8024372242**

Option 3 ID : **8024372243**

Option 4 ID : **8024372244**

Status : **Answered**

Chosen Option : **2,3**

Q.4 Which of the following sets are in bijection with \mathbb{R} ?

Options

1. Set of all maps from $\{0,1\}$ to \mathbb{N}
2. Set of all maps from \mathbb{N} to $\{0,1\}$
3. Set of all subsets of \mathbb{N}
4. Set of all subsets of \mathbb{R}

Question Type : **MSQ**

Question ID : **802437551**

Option 1 ID : **8024372201**

Option 2 ID : **8024372202**

Option 3 ID : **8024372203**

Option 4 ID : **8024372204**

Status : **Answered**

Chosen Option : **2,3,4**

Q.5 Let n be a positive integer and F be a non-empty proper subset of $\{1,2, \dots, n\}$. Define

$$\langle x, y \rangle_F = \sum_{k \in F} x_k y_k, \quad x = (x_1, \dots, x_n), \quad y = (y_1, \dots, y_n) \in \mathbb{R}^n.$$

Let $T = \{x \in \mathbb{R}^n : \langle x, x \rangle_F = 0\}$. Which of the following statements are true?

For $y \in \mathbb{R}^n, y \neq 0$

Options

1. $\inf_{x \in T} \langle x + y, x + y \rangle_F = \langle y, y \rangle_F$
2. $\sup_{x \in T} \langle x + y, x + y \rangle_F = \langle y, y \rangle_F$
3. $\inf_{x \in T} \langle x + y, x + y \rangle_F < \langle y, y \rangle_F$
4. $\sup_{x \in T} \langle x + y, x + y \rangle_F > \langle y, y \rangle_F$

Question Type : **MSQ**

Question ID : **802437565**

Option 1 ID : **8024372257**

Option 2 ID : **8024372258**

Option 3 ID : **8024372259**

Option 4 ID : **8024372260**

Status : **Not Attempted and
Marked For Review**

Chosen Option : **--**

Q.6

Which of the following statements are true?

Options

1. The series $\sum_{n \geq 1} \frac{(-1)^n}{\sqrt{n}}$ is convergent
2. The series $\sum_{n \geq 1} \frac{(-1)^n}{\sqrt{n} + n}$ is absolutely convergent
3. The series $\sum_{n \geq 1} \frac{[1 + (-1)^n]\sqrt{n} + \log n}{n^{3/2}}$ is convergent
4. The series $\sum_{n \geq 1} \frac{((-1)^n \sqrt{n} + 1)}{n^{3/2}}$ is convergent

Question Type : MSQ

Question ID : 802437552

Option 1 ID : 8024372205

Option 2 ID : 8024372206

Option 3 ID : 8024372207

Option 4 ID : 8024372208

Status : Answered

Chosen Option : 1,3,4

Q.7

Define

$$f(x, y) = \begin{cases} \frac{x^2 - y^2}{x^2 + y^2} & \text{for } (x, y) \neq (0, 0) \\ 0 & \text{for } (x, y) = (0, 0) \end{cases}$$

Which of the following statements are true?

Options

1. f is continuous at $(0, 0)$
2. f is bounded in a neighbourhood of $(0, 0)$
3. f is not bounded in any neighbourhood of $(0, 0)$
4. f has all directional derivatives at $(0, 0)$

Question Type : MSQ

Question ID : 802437559

Option 1 ID : 8024372233

Option 2 ID : 8024372234

Option 3 ID : 8024372235

Option 4 ID : 8024372236

Status : Answered

Chosen Option : 2,4

Q.8

Define

$$f(x, y) = \begin{cases} \frac{x^3}{x^2+y^2} & \text{for } (x, y) \neq (0,0) \\ 0 & \text{for } (x, y) = (0,0) \end{cases}$$

Which of the following statements are true?

Options

1. f is discontinuous at $(0, 0)$
2. f is continuous at $(0, 0)$
3. all directional derivatives of f at $(0, 0)$ exist
4. f is not differentiable at $(0, 0)$

Question Type : MSQ

Question ID : 802437558

Option 1 ID : 8024372229

Option 2 ID : 8024372230

Option 3 ID : 8024372231

Option 4 ID : 8024372232

Status : Answered

Chosen Option : 2

Q.9

Let A be an $n \times n$ matrix such that the first 3 rows of A are linearly independent and the first 5 columns of A are linearly independent. Which of the following statements are true?

Options

1. A has at least 5 linearly independent rows
2. $3 \leq \text{rank } A \leq 5$
3. $\text{rank } A \geq 5$
4. $\text{rank } A^2 \geq 5$

Question Type : MSQ

Question ID : 802437564

Option 1 ID : 8024372253

Option 2 ID : 8024372254

Option 3 ID : 8024372255

Option 4 ID : 8024372256

Status : Answered

Chosen Option : 2,4

- Q.10** Let $v \in \mathbb{R}^3$ be a non-zero vector. Define a linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by $T(x) = x - 2 \frac{x \cdot v}{v \cdot v} v$, where $x \cdot y$ denotes the standard inner product in \mathbb{R}^3 .
- Which of the following statements are true?

Options

1. The eigenvalues of T are $+1, -1$
2. The determinant of T is -1
3. The trace of T is $+1$
4. T is distance preserving

Question Type : **MSQ**
 Question ID : **802437566**
 Option 1 ID : **8024372261**
 Option 2 ID : **8024372262**
 Option 3 ID : **8024372263**
 Option 4 ID : **8024372264**
 Status : **Answered**
 Chosen Option : **1,2,4**

- Q.11** Consider two series

$$A(x) = \sum_{n=0}^{\infty} x^n(1-x) \text{ and } B(x) = \sum_{n=0}^{\infty} (-1)^n x^n(1-x)$$

where $x \in [0,1]$.

Which of the following statements are true?

Options

1. Both $A(x)$ and $B(x)$ converge pointwise
2. Both $A(x)$ and $B(x)$ converge uniformly
3. $A(x)$ converges uniformly but $B(x)$ does not
4. $B(x)$ converges uniformly but $A(x)$ does not

Question Type : **MSQ**
 Question ID : **802437554**
 Option 1 ID : **8024372213**
 Option 2 ID : **8024372214**
 Option 3 ID : **8024372215**
 Option 4 ID : **8024372216**
 Status : **Answered**
 Chosen Option : **1,2**

Q.12 A quadratic form $Q(x, y, z)$ over \mathbb{R} represents 0 non trivially if there exists $(a, b, c) \in \mathbb{R}^3 \setminus \{(0, 0, 0)\}$ such that $Q(a, b, c) = 0$. Which of the following quadratic forms $Q(x, y, z)$ over \mathbb{R} represent 0 non trivially?

Options

1. $Q(x, y, z) = xy + z^2$
2. $Q(x, y, z) = x^2 + 3y^2 - 2z^2$
3. $Q(x, y, z) = x^2 - xy + y^2 + z^2$
4. $Q(x, y, z) = x^2 + xy + z^2$

Question Type : **MSQ**

Question ID : **802437567**

Option 1 ID : **8024372265**

Option 2 ID : **8024372266**

Option 3 ID : **8024372267**

Option 4 ID : **8024372268**

Status : **Answered**

Chosen Option : 1,4

Q.13 Suppose that $\{f_n\}$ is a sequence of real-valued functions on \mathbb{R} . Suppose it converges to a continuous function f uniformly on each closed and bounded subset of \mathbb{R} . Which of the following statements are true?

Options 1.

1. The sequence $\{f_n\}$ converges to f uniformly on \mathbb{R}
2. The sequence $\{f_n\}$ converges to f pointwise on \mathbb{R}
3. For all sufficiently large n , the function f_n is bounded
4. For all sufficiently large n the function f_n is continuous

Question Type : **MSQ**

Question ID : **802437556**

Option 1 ID : **8024372221**

Option 2 ID : **8024372222**

Option 3 ID : **8024372223**

Option 4 ID : **8024372224**

Status : **Not Answered**

Chosen Option : --

Q.14

For $p \in \mathbb{R}$, consider the improper integral

$$I_p = \int_0^1 t^p \sin t \, dt.$$

Which of the following statements are true?

Options

1. I_p is convergent for $p = -1/2$
2. I_p is divergent for $p = -3/2$
3. I_p is convergent for $p = 4/3$
4. I_p is divergent for $p = -4/3$

Question Type : **MSQ**Question ID : **802437555**Option 1 ID : **8024372217**Option 2 ID : **8024372218**Option 3 ID : **8024372219**Option 4 ID : **8024372220**Status : **Not Answered**

Chosen Option : --

Q.15

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $f(x, y) = \begin{cases} \frac{2xy}{x^2+y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0). \end{cases}$

Define $g(x, y) = \sum_{n=1}^{\infty} \frac{f((x-n), (y-n))}{2^n}$.

Which of the following statements are true?

Options 1.

1. The function $h(y) = g(c, y)$ is continuous on \mathbb{R} for all c
2. g is continuous from \mathbb{R}^2 into \mathbb{R}
3. g is not a well-defined function
4. g is continuous on $\mathbb{R}^2 \setminus \{(k, k)\}_{k \in \mathbb{N}}$

Question Type : MSQ

Question ID : 802437553

Option 1 ID : 8024372209

Option 2 ID : 8024372210

Option 3 ID : 8024372211

Option 4 ID : 8024372212

Status : Not Answered

Chosen Option : --

Q.16

Let A and B be $n \times n$ real matrices and let $C = \begin{pmatrix} A & B \\ B & A \end{pmatrix}$.

Which of the following statements are true?

Options 1.

1. If λ is an eigenvalue of $A + B$ then λ is an eigenvalue of C
2. If λ is an eigenvalue of $A - B$ then λ is an eigenvalue of C
3. If λ is an eigenvalue of A or B then λ is an eigenvalue of C
4. All eigenvalues of C are real

Question Type : MSQ

Question ID : 802437562

Option 1 ID : 8024372245

Option 2 ID : 8024372246

Option 3 ID : 8024372247

Option 4 ID : 8024372248

Status : Answered

Chosen Option : 1,2,3

Q.17 Let $f(x) = e^{-x}$ and $g(x) = e^{-x^2}$. Which of the following statements are true?

Options

1. Both f and g are uniformly continuous on \mathbb{R}
2. f is uniformly continuous on every interval of the form $[a, +\infty)$, $a \in \mathbb{R}$
3. g is uniformly continuous on \mathbb{R}
4. $f(x)g(x)$ is uniformly continuous on \mathbb{R}

Question Type : **MSQ**
 Question ID : **802437557**
 Option 1 ID : **8024372225**
 Option 2 ID : **8024372226**
 Option 3 ID : **8024372227**
 Option 4 ID : **8024372228**
 Status : **Answered**
 Chosen Option : **2,3,4**

Q.18 Let A be an $n \times n$ real matrix. Let b be an $n \times 1$ vector. Suppose $Ax = b$ has no solution.

Which of the following statements are true?

Options

1. There exists an $n \times 1$ vector c such that $Ax = c$ has a unique solution
2. There exist infinitely many vectors c such that $Ax = c$ has no solution
3. If y is the first column of A then $Ax = y$ has a unique solution
4. $\det A = 0$

Question Type : **MSQ**
 Question ID : **802437563**
 Option 1 ID : **8024372249**
 Option 2 ID : **8024372250**
 Option 3 ID : **8024372251**
 Option 4 ID : **8024372252**
 Status : **Answered**
 Chosen Option : **2,4**

Q.19

For $z \neq -i$, let $f(z) = \exp\left(\frac{1}{z+i}\right) - 1$. Which of the following are true?

Options

1. f has finitely many zeros
2. f has a sequence of zeros that converges to a removable singularity of f
3. f has a sequence of zeros that converges to a pole of f
4. f has a sequence of zeros that converges to an essential singularity of f

Question Type : MSQ

Question ID : 802437569

Option 1 ID : 8024372273

Option 2 ID : 8024372274

Option 3 ID : 8024372275

Option 4 ID : 8024372276

Status : Not Answered

Chosen Option : --

Q.20

Let G be a finite group. Which of the following are true?

Options

1. If $g \in G$ has order m and if $n \geq 1$ divides m , then G has a subgroup of order n .
2. If for any two subgroups A and B of G , either $A \subset B$ or $B \subset A$, then G is cyclic.
3. If G is cyclic, then for any two subgroups A and B of G , either $A \subset B$ or $B \subset A$.
4. If for every positive integer m dividing $|G|$, G has a subgroup of order m , then G is abelian.

Question Type : MSQ

Question ID : 802437575

Option 1 ID : 8024372297

Option 2 ID : 8024372298

Option 3 ID : 8024372299

Option 4 ID : 8024372300

Status : Answered

Chosen Option : 3,4

Q.21

Let $SL_2(\mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Z}) : ad - bc = 1 \right\}$ and for any prime p , let

$$\Gamma(p) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z}) \mid \begin{matrix} a \equiv 1 \pmod{p}, d \equiv 1 \pmod{p} \\ c \equiv 0 \pmod{p}, b \equiv 0 \pmod{p} \end{matrix} \right\}$$

Which of the following are true?

Options

1. $\Gamma(p)$ is a subgroup of $SL_2(\mathbb{Z})$
2. $\Gamma(p)$ is not a normal subgroup of $SL_2(\mathbb{Z})$
3. $\Gamma(p)$ has atleast two elements
4. $\Gamma(p)$ is uncountable

Question Type : MSQ

Question ID : 802437574

Option 1 ID : 8024372293

Option 2 ID : 8024372294

Option 3 ID : 8024372295

Option 4 ID : 8024372296

Status : Not Answered

Chosen Option : --

Q.22

Which of the following statements are true?

Options

1. \mathbb{Q} has countably many subgroups
2. \mathbb{Q} has uncountably many subsets
- 3.

Every finitely generated subgroup of \mathbb{Q} is cyclic

4. \mathbb{Q} is isomorphic to $\mathbb{Q} \times \mathbb{Q}$ as groups

Question Type : MSQ

Question ID : 802437573

Option 1 ID : 8024372289

Option 2 ID : 8024372290

Option 3 ID : 8024372291

Option 4 ID : 8024372292

Status : Answered

Chosen Option : 2,3

Q.23 Let f be a holomorphic function on the open unit disc $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$.

Suppose that $|f| \geq 1$ on \mathbb{D} and $f(0) = i$.

Which of the following are possible values of $f\left(\frac{1}{2}\right)$?

- Options**
1. $-i$
 2. i
 3. 1
 4. -1

Question Type : **MSQ**

Question ID : **802437570**

Option 1 ID : **8024372277**

Option 2 ID : **8024372278**

Option 3 ID : **8024372279**

Option 4 ID : **8024372280**

Status : **Answered**

Chosen Option : 1,3

Q.24 Let R, S be commutative rings with unity, $f: R \rightarrow S$ be a surjective ring homomorphism,

$Q \subseteq S$ be a non-zero prime ideal. Which of the following statements are true?

- Options**
1. $f^{-1}(Q)$ is a non-zero prime ideal in R
 2. $f^{-1}(Q)$ is a maximal ideal in R if R is a *PID*
 3. $f^{-1}(Q)$ is a maximal ideal in R if R is a finite commutative ring with unity
 4. $f^{-1}(Q)$ is a maximal ideal in R if $x^5 = x$ for all $x \in R$

Question Type : **MSQ**

Question ID : **802437576**

Option 1 ID : **8024372301**

Option 2 ID : **8024372302**

Option 3 ID : **8024372303**

Option 4 ID : **8024372304**

Status : **Answered**

Chosen Option : 1,3

Q.25 Let X be a non-empty set. Suppose that τ_1 and τ_2 are two topologies over X , such that $\tau_2 \subset \tau_1$.

Which of the following statements imply that $\tau_1 = \tau_2$?

Options

1. (X, τ_1) is compact and τ_1 is T_2 (Hausdorff)
2. (X, τ_1) is compact and τ_2 is T_2 (Hausdorff)
3. The connected components of both (X, τ_1) and (X, τ_2) are same
4. For any subset $A \subset X$ the closure of A in (X, τ_2) is contained in the closure of A in (X, τ_1)

Question Type : **MSQ**

Question ID : **802437580**

Option 1 ID : **8024372317**

Option 2 ID : **8024372318**

Option 3 ID : **8024372319**

Option 4 ID : **8024372320**

Status : **Not Answered**

Chosen Option : **--**

Q.26 Consider the polynomial $f(x) = x^2 + 3x - 1$. Which of the following statements are true?

Options

1. f is irreducible over $\mathbb{Z}[\sqrt{13}]$
2. f is irreducible over \mathbb{Q}
3. f is reducible over $\mathbb{Q}[\sqrt{13}]$
4. $\mathbb{Z}[\sqrt{13}]$ is a unique factorization domain

Question Type : **MSQ**

Question ID : **802437577**

Option 1 ID : **8024372305**

Option 2 ID : **8024372306**

Option 3 ID : **8024372307**

Option 4 ID : **8024372308**

Status : **Answered**

Chosen Option : **2,4**

Q.27 Let n be a positive integer. For a real number

$$R > 1, \text{ let } z(\theta) = Re^{i\theta}, 0 \leq \theta < 2\pi.$$

The set $\{\theta \in [0, 2\pi) : |z(\theta)^n + 1| = |z(\theta)|^n - 1\}$ contains which of the following sets?

Options

1. $\{\theta \in [0, 2\pi) : \cos n\theta = 1\}$
2. $\{\theta \in [0, 2\pi) : \sin n\theta = 1\}$
3. $\{\theta \in [0, 2\pi) : \cos n\theta = -1\}$
4. $\{\theta \in [0, 2\pi) : \sin n\theta = -1\}$

Question Type : **MSQ**

Question ID : **802437572**

Option 1 ID : **8024372285**

Option 2 ID : **8024372286**

Option 3 ID : **8024372287**

Option 4 ID : **8024372288**

Status : **Not Answered**

Chosen Option : **--**

Q.28 Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ be the open unit disc and let $f: \mathbb{D} \rightarrow \mathbb{D}$ be a holomorphic function. Suppose that $f(0) = 0$ and $f'(0) = 0$. Which of the following are possible values of $f\left(\frac{1}{2}\right)$?

Options

1. $1/4$
2. $-1/4$
3. $1/3$
4. $-1/3$

Question Type : **MSQ**

Question ID : **802437571**

Option 1 ID : **8024372281**

Option 2 ID : **8024372282**

Option 3 ID : **8024372283**

Option 4 ID : **8024372284**

Status : **Answered**

Chosen Option : **2**

Q.29 Let p be an odd prime such that $p \equiv 2 \pmod{3}$. Let \mathbb{F}_p be the field with p elements. Consider the subset E of $\mathbb{F}_p \times \mathbb{F}_p$ given by

$$E = \{(x, y) \in \mathbb{F}_p \times \mathbb{F}_p : y^2 = x^3 + 1\}.$$

Which of the following are true?

- Options
1. E has atleast two elements
 2. E has atleast $2p$ elements
 3. E can have p^2 elements
 4. E has atleast $2p$ elements

Question Type : **MSQ**

Question ID : **802437578**

Option 1 ID : **8024372309**

Option 2 ID : **8024372310**

Option 3 ID : **8024372311**

Option 4 ID : **8024372312**

Status : **Not Answered**

Chosen Option : --

Q.30 Consider the subset of \mathbb{R}^2 defined as follows:

$$A = \{(x, y) \in \mathbb{R} \times \mathbb{R} : (x - 1)(x - 2)(y - 3)(y + 4) = 0\}$$

Which of the following statements are true?

- Options
1. A is connected
 2. A is compact
 3. A is closed
 4. A is dense

Question Type : **MSQ**

Question ID : **802437579**

Option 1 ID : **8024372313**

Option 2 ID : **8024372314**

Option 3 ID : **8024372315**

Option 4 ID : **8024372316**

Status : **Not Answered**

Chosen Option : --

Q.31 The following two-point boundary value problem

$$\begin{cases} y''(x) + \lambda y(x) = 0 & \text{for } x \in (0, \pi) \\ y(0) = 0 \\ y(\pi) = 0 \end{cases}$$

has a trivial solution $y = 0$. It also has a non-trivial solution for

Options

1. no values of $\lambda \in \mathbb{R}$
2. $\lambda = 1$
3. $\lambda = n^2$ for all $n \in \mathbb{N}, n > 1$
4. $\lambda \leq 0$

Question Type : **MSQ**

Question ID : **802437581**

Option 1 ID : **8024372321**

Option 2 ID : **8024372322**

Option 3 ID : **8024372323**

Option 4 ID : **8024372324**

Status : **Not Answered**

Chosen Option : --

Q.32 Consider a dynamical system with the Lagrangian function $L(q, \dot{q}) = T - U$, where the kinetic energy

$$T = a(q)\dot{q}^2 \geq 0$$

and the potential energy $U := U(q)$ with $a(q) > 0$. Which of the following statements are true?

Options

The associated Lagrange's equation is

1.
$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}} = \frac{\partial L}{\partial q}$$

The associated Lagrange's equation is

2.
$$\frac{d}{dt} \frac{\partial L}{\partial q} = \frac{\partial L}{\partial \dot{q}}$$

3.

The point (q_0, \dot{q}_0) is an equilibrium position of the dynamical system if and only if

$$\dot{q}_0 = 0 \quad \text{and} \quad \left. \frac{\partial U}{\partial q} \right|_{q=q_0} = 0$$

4.

The point (q_0, \dot{q}_0) is an equilibrium position of the dynamical system if and only if

$$\dot{q}_0 = 0 \quad \text{and} \quad \left. \frac{\partial U}{\partial q} \right|_{q=q_0} > 0$$

Question Type : **MSQ**

Question ID : **802437592**

Option 1 ID : **8024372365**

Option 2 ID : **8024372366**

Option 3 ID : **8024372367**

Option 4 ID : **8024372368**

Status : **Not Answered**

Chosen Option : --

Q.33 Consider the Cauchy problem

$$\begin{cases} \frac{\partial^2 u}{\partial x \partial y} = 0, |x| < 1, 0 < y < 1 \\ u(x, x^2) = 0, \quad \frac{\partial u}{\partial y}(x, x^2) = g(x), |x| < 1. \end{cases}$$

Which of the following statements are true?

Options 1.

1. A necessary condition for a solution to exist is that g is an odd function

2.

A necessary condition for a solution to exist is that g is an even function

3.

The solution (if it exists) is given by $u(x, y) = 2 \int_x^{\sqrt{y}} z g(z) dz$

4.

The solution (if it exists) is given by $u(x, y) = 2 \int_{\sqrt{y}}^{x^2} z g(z) dz$

Question Type : **MSQ**

Question ID : **802437585**

Option 1 ID : **8024372337**

Option 2 ID : **8024372338**

Option 3 ID : **8024372339**

Option 4 ID : **8024372340**

Status : **Not Answered**

Chosen Option : --

Q.34 Let A be an $n \times n$ matrix with distinct eigenvalues $\{\lambda_1, \dots, \lambda_n\}$ with corresponding linearly independent eigenvectors $\{v_1, \dots, v_n\}$.

Then, the non-homogeneous differential equation

$$x'(t) = Ax(t) + e^{\lambda_1 t} v_1$$

Options 1.

1. does not have a solution of the form $e^{\lambda_1 t} a$ for any vector $a \in \mathbb{R}^n$

2.

has a solution of the form $e^{\lambda_1 t} a$ for some vector $a \in \mathbb{R}^n$

3.

has a solution of the form $e^{\lambda_1 t} a + t e^{\lambda_1 t} b$ for some vectors $a, b \in \mathbb{R}^n$

4.

does not have a solution of the form $e^{\lambda_1 t} a + t e^{\lambda_1 t} b$ for any vectors $a, b \in \mathbb{R}^n$

Question Type : **MSQ**

Question ID : **802437582**

Option 1 ID : **8024372325**

Option 2 ID : **8024372326**

Option 3 ID : **8024372327**

Option 4 ID : **8024372328**

Status : **Not Answered**

Chosen Option : --

Q.35

For the Fredholm integral equation

$$y(s) = \lambda \int_0^1 e^s e^t y(t) dt$$

Which of the following statements are true?

Options 1.

1. It has a non-trivial solution satisfying $\int_0^1 e^t y(t) dt = 0$
2. Only the trivial solution satisfies $\int_0^1 e^t y(t) dt = 0$
3. It has non-trivial solution for all $\lambda \neq 0$
4. It has non-trivial solutions only if $\lambda = \frac{2}{e^2-1}$ and $\int_0^1 e^t y(t) dt \neq 0$

Question Type : MSQ

Question ID : 802437590

Option 1 ID : 8024372357

Option 2 ID : 8024372358

Option 3 ID : 8024372359

Option 4 ID : 8024372360

Status : Answered

Chosen Option : 2,4

Q.36

Consider the functional

$$J(y) = \int_0^\pi ((y')^2 - ky^2) dx \text{ with boundary conditions } y(0) = 0, y(\pi) = 0$$

Which of the following statements are true?

Options

1. It has a unique extremal for all $k \in \mathbb{R}$
2. It has at most one extremal if \sqrt{k} is not an integer
3. It has infinitely many extremals if \sqrt{k} is an integer
4. It has a unique extremal if \sqrt{k} is an integer

Question Type : MSQ

Question ID : 802437589

Option 1 ID : 8024372353

Option 2 ID : 8024372354

Option 3 ID : 8024372355

Option 4 ID : 8024372356

Status : Not Answered

Chosen Option : --

Q.37

Consider the solutions

$$y_1 := \begin{pmatrix} e^{-3t} \\ e^{-3t} \\ 0 \end{pmatrix} \text{ and } y_2 := \begin{pmatrix} 0 \\ e^{-5t} \\ e^{-5t} \end{pmatrix}$$

to the homogeneous linear system of differential equation

$$(*) \quad y'(t) = \begin{pmatrix} -5 & 2 & -2 \\ 1 & -4 & -1 \\ -1 & 1 & -6 \end{pmatrix} y(t).$$

Which of the following statements are true?

Options 1.

1. y_1 and y_2 form a basis for the set of all solutions to (*)
2. y_1 and y_2 are linearly independent but do not form a basis for the set of all solutions to (*)
3. There exists another solution y_3 such that $\{y_1, y_2, y_3\}$ form a basis for the set of all solutions to (*)
4. y_1 and y_2 are linearly dependent

Question Type : **MSQ**Question ID : **802437583**Option 1 ID : **8024372329**Option 2 ID : **8024372330**Option 3 ID : **8024372331**Option 4 ID : **8024372332**Status : **Not Answered**Chosen Option : **--**

Q.38 Fix a $\alpha \in (0, 1)$. Consider the iteration defined by

$$(*) \quad x_{k+1} = \frac{1}{2}(x_k^2 + \alpha), \quad k = 0, 1, 2, \dots$$

The above iteration has two distinct fixed points ζ_1 and ζ_2 such that

$$0 < \zeta_1 < 1 < \zeta_2.$$

Which of the following statements are true?

Options 1.

The iteration (*) is equivalent to the recurrence relation

$$x_{k+1} - \zeta_1 = \frac{1}{2}(x_k + \zeta_1)(x_k - \zeta_1), \quad k = 0, 1, 2, \dots$$

2.

The iteration (*) is equivalent to the recurrence relation

$$x_{k+1} - \zeta_1 = \frac{1}{2}(x_k + \zeta_2)(x_k - \zeta_1), \quad k = 0, 1, 2, \dots$$

3. If $0 \leq x_0 < \zeta_2$ then $\lim_{k \rightarrow \infty} x_k = \zeta_1$

4. If $-\zeta_2 < x_0 \leq 0$ then $\lim_{k \rightarrow \infty} x_k = \zeta_1$

Question Type : MSQ

Question ID : 802437586

Option 1 ID : 8024372341

Option 2 ID : 8024372342

Option 3 ID : 8024372343

Option 4 ID : 8024372344

Status : Not Answered

Chosen Option : --

Q.39

Consider the partial differential equation

$$z = x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} + \frac{\partial z}{\partial x} \frac{\partial z}{\partial y}$$

Which of the following statements are true?

Options 1.

1. The complete integral is $z = xa + yb + ab$, a, b arbitrary constants

2.

The complete integral is $z = xa + yb + \sqrt{a^2 + b^2}$, a, b arbitrary constants

3.

The particular solution passing through $x = 0$ and $z = y^2$ is $\left(\frac{x}{4} - y\right)^2$

4.

The particular solution passing through $x = 0$ and $z = y^2$ is $\left(\frac{x}{4} + y\right)^2$ Question Type : **MSQ**Question ID : **802437591**Option 1 ID : **8024372361**Option 2 ID : **8024372362**Option 3 ID : **8024372363**Option 4 ID : **8024372364**Status : **Not Answered**Chosen Option : **--**

Q.40 The extremal of the functional

$$J(y) = \int_0^1 e^x \sqrt{1 + (y')^2} dx, y \in C^2[0, 1]$$

is of the form

Options 1.

1. $y = \sec^{-1}\left(\frac{x}{c_1}\right) + c_2$, where c_1 and c_2 are arbitrary constants

2.

2. $y = \sec^{-1}\left(\frac{x}{c_1}\right) + c_2$, where $|c_1| < 1$ and c_2 is an arbitrary constant

3.

3. $y = \tan^{-1}\left(\frac{x}{c_1}\right) + c_2$, where c_1 and c_2 are arbitrary constants

4.

4. $y = \tan^{-1}\left(\frac{x}{c_1}\right) + c_2$, where $|c_1| > 1$ and c_2 is an arbitrary constant

Question Type : **MSQ**

Question ID : **802437588**

Option 1 ID : **8024372349**

Option 2 ID : **8024372350**

Option 3 ID : **8024372351**

Option 4 ID : **8024372352**

Status : **Not Answered**

Chosen Option : --

Q.41 Consider the function $f: [0, 1] \rightarrow \mathbb{R}$ defined by

$$f(x) := \begin{cases} 2^{-\left\{1 + \left(\log_2\left(\frac{1}{x}\right)\right)^{\frac{1}{\beta}}\right\}} & \text{for } x \in (0, 1] \\ 0 & \text{for } x = 0, \end{cases}$$

where $\beta \in (0, \infty)$ is a parameter. Consider the iterations

$$x_{k+1} = f(x_k) \quad , \quad k = 0, 1, \dots; x_0 > 0.$$

Which of the following statements are true about the iteration ?

Options 1.

For $\beta = 1$, the sequence $\{x_k\}$ converges to 0 linearly with asymptotic rate of convergence $\log_{10} 2$

2.

For $\beta > 1$, the sequence $\{x_k\}$ does not converge to 0

3.

For $\beta \in (0, 1)$, the sequence $\{x_k\}$ converges to 0 sublinearly

4.

For $\beta \in (0, 1)$, the sequence $\{x_k\}$ converges to 0 superlinearly

Question Type : **MSQ**

Question ID : **802437587**

Option 1 ID : **8024372345**

Option 2 ID : **8024372346**

Option 3 ID : **8024372347**

Option 4 ID : **8024372348**

Status : **Not Answered**

Chosen Option : --

Q.42

Consider the partial differential equations

$$(i) \quad \frac{\partial^2 u}{\partial x^2} + 2 \frac{\partial^2 u}{\partial x \partial y} + (1 - \operatorname{sgn}(y)) \frac{\partial^2 u}{\partial y^2} = 0$$

$$(ii) \quad y \frac{\partial^2 u}{\partial x^2} + x \frac{\partial^2 u}{\partial y^2} = 0$$

Which of the following statements are true?

Options 1.

Equation (i) is parabolic for $y > 0$ and elliptic for $y < 0$

2.

Equation (i) is hyperbolic for $y > 0$ and elliptic for $y < 0$

3.

Equation (ii) is elliptic in I and III quadrant and hyperbolic in II and IV quadrant

4.

Equation (ii) is hyperbolic in I and III quadrant and elliptic in II and IV quadrant

Question Type : MSQ

Question ID : 802437584

Option 1 ID : 8024372333

Option 2 ID : 8024372334

Option 3 ID : 8024372335

Option 4 ID : 8024372336

Status : Not Answered

Chosen Option : --

Q.43

Let $p > 1$ and $1 > \rho \geq 0$. Consider a multiple linear regression problem with p independent variables X_1, X_2, \dots, X_p and a dependent variable Y . Suppose that the correlation between Y and X_i is ρ and the correlation between X_i and X_j is also ρ for all $1 \leq i < j \leq p$. Which of the following are correct?

Options 1.

The multiple correlation between Y and (X_1, \dots, X_p) is larger than or equal to ρ

2.

The multiple correlation between Y and (X_1, \dots, X_p) will be ρ if $\rho = 0$

3.

The multiple correlation between Y and (X_1, \dots, X_p) will be ρ only if $\rho = 0$

4.

The multiple correlation between Y and (X_1, \dots, X_p) tends to 1 as $p \rightarrow \infty$

Question Type : MSQ

Question ID : 802437604

Option 1 ID : 8024372413

Option 2 ID : 8024372414

Option 3 ID : 8024372415

Option 4 ID : 8024372416

Status : Not Answered

Chosen Option : --

Q.44 Consider a M/M/1 queue with arrival rate λ and service rate μ . Let $Q_0 = 0$ and Q_t denote the queue length at time t . Which of the following statements are true?

Options 1.

(Q_t) admits a stationary distribution if and only if $\lambda \leq \mu$

2.

The stationary distribution of the process (Q_t) is geometric, when it exists

3. $\lim_{t \rightarrow \infty} P(Q_t > k) = 1$ for all $k < \infty$ if $\lambda > \mu$

4. $\lim_{t \rightarrow \infty} P(Q_t > k) = 2^{-(k+1)}$ for all $k < \infty$ if $\lambda = \frac{\mu}{2}$

Question Type : **MSQ**

Question ID : **802437610**

Option 1 ID : **8024372437**

Option 2 ID : **8024372438**

Option 3 ID : **8024372439**

Option 4 ID : **8024372440**

Status : **Not Answered**

Chosen Option : --

Q.45 X has binomial distribution with parameters n and p . Suppose that n is given and the unknown parameter p has prior distribution, which is uniform on the interval $[0, 1]$. Consider the squared error loss function and the observation $X = n$. Identify the correct statement.

Options

1. The Bayes estimate of p is $\left(\frac{n+1}{n+2}\right)$

2. The Bayes estimate of p is $2^{-1/(n+1)}$

3.

The median of the posterior distribution of p is $2^{-1/(n+1)}$

4.

The median of the posterior distribution of p is $\left(\frac{n+1}{n+2}\right)$

Question Type : **MSQ**

Question ID : **802437601**

Option 1 ID : **8024372401**

Option 2 ID : **8024372402**

Option 3 ID : **8024372403**

Option 4 ID : **8024372404**

Status : **Not Answered**

Chosen Option : --

Q.46 Consider a Markov chain with a countable state space S . Identify the correct statements.

Options 1.

If the Markov chain is aperiodic and irreducible then there exists a stationary distribution

2.

If the Markov chain is aperiodic and irreducible then there is at most one stationary distribution

3.

If S is finite then there exists a stationary distribution

4.

If S is finite then there is exactly one stationary distribution

Question Type : **MSQ**

Question ID : **802437595**

Option 1 ID : **8024372377**

Option 2 ID : **8024372378**

Option 3 ID : **8024372379**

Option 4 ID : **8024372380**

Status : **Not Answered**

Chosen Option : --

Q.47 Let X_1, X_2, \dots, X_n be i.i.d. with the common probability mass function

$$p(x|\theta) = \theta^x(1-\theta)^{1-x}, \quad x = 0 \text{ or } 1, \text{ and } 0 \leq \theta \leq \frac{1}{2}$$

Then

Options 1.

the method of moments estimator of θ is $\frac{1}{2n} \sum_{i=1}^n X_i$

2. the MLE of θ is $\min_{1 \leq i \leq n} X_i$

3.

the method of moments estimator of θ is $\min_{1 \leq i \leq n} X_i$

4. the MLE of θ is $\min \left\{ \frac{1}{n} \sum_{i=1}^n X_i, \frac{1}{2} \right\}$

Question Type : **MSQ**

Question ID : **802437599**

Option 1 ID : **8024372393**

Option 2 ID : **8024372394**

Option 3 ID : **8024372395**

Option 4 ID : **8024372396**

Status : **Not Answered**

Chosen Option : --

- Q.48** Let $n > 2$ and $0 < \theta < \frac{\pi}{2}$ be fixed. Let X_1, \dots, X_n be i.i.d. normal random variables with mean zero and variance $\sigma^2 > 0$. For $i = 1, \dots, n$ define
- $$Y_{2i-1} = X_i \cos \theta \text{ and } Y_{2i} = X_i \sin \theta. \text{ Further, let } Z^T = (Y_1, Y_2, \dots, Y_{2n}), \text{ and}$$
- $$V^T = (X_1, Y_1, Y_2, X_2, Y_3, Y_4, \dots, X_n, Y_{2n-1}, Y_{2n}).$$
- Which of the following statements are correct?

Options

1. Z^T has a multivariate normal distribution
2. There exists a constant C , such that $CZ^T Z$ has a chi-square distribution
3. V^T has a multivariate normal distribution
4. $E\left(\frac{1}{V^T V}\right) < \infty$

Question Type : **MSQ**
 Question ID : **802437605**
 Option 1 ID : **8024372417**
 Option 2 ID : **8024372418**
 Option 3 ID : **8024372419**
 Option 4 ID : **8024372420**
 Status : **Not Answered**
 Chosen Option : --

- Q.49** Let $X_i = \theta + \varepsilon_i, 1 \leq i \leq n$ where $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ are i.i.d. with pdf $g(\varepsilon) = |\varepsilon|, -1 < \varepsilon < 1$. Let
- $$T_1 = \frac{1}{n} \sum_{i=1}^n X_i \text{ and } T_2 = X_{\left(\left[\frac{3n}{4}\right]+1\right)}, \text{ the sample 75th percentile. Which of the following are correct?}$$

Options 1.

1. T_1 is a consistent and asymptotically normal estimator of θ
2. $T_2 - \frac{1}{\sqrt{2}}$ is a consistent and asymptotically normal estimator of θ
3. The asymptotic variance of T_1 is $\frac{1}{2n}$
4. The asymptotic variance of T_2 is $\frac{3}{8n}$

Question Type : **MSQ**
 Question ID : **802437609**
 Option 1 ID : **8024372433**
 Option 2 ID : **8024372434**
 Option 3 ID : **8024372435**
 Option 4 ID : **8024372436**
 Status : **Not Answered**
 Chosen Option : --

Q.50 Suppose $\lambda(t)$ for $t \geq 0$ is a continuous hazard function of a non-negative random variable X , where $\lambda(t) \geq 1$. Which of the following statements are always true?

Options

1. $\frac{1}{\lambda(t)}$ is a hazard function
2. $\lambda^2(t)$ is also a hazard function
3. $c\lambda(t)$ for $c \geq 0$ is also a hazard function
4. $\log \lambda(t)$ is a hazard function

Question Type : **MSQ**

Question ID : **802437608**

Option 1 ID : **8024372429**

Option 2 ID : **8024372430**

Option 3 ID : **8024372431**

Option 4 ID : **8024372432**

Status : **Not Answered**

Chosen Option : --

Q.51 In a Randomized Block Design with one observation per cell, and data satisfying the standard linear model, which of the following are correct?

Options

1. Mean treatment effects are estimable
2. Mean block effects are estimable
3. Treatment-Block interactions are NOT estimable
4. Treatment and block effects as well as treatment-block interactions are estimable

Question Type : **MSQ**

Question ID : **802437607**

Option 1 ID : **8024372425**

Option 2 ID : **8024372426**

Option 3 ID : **8024372427**

Option 4 ID : **8024372428**

Status : **Not Answered**

Chosen Option : --

Q.52 Suppose X_1, X_2, \dots, X_n are i.i.d. random variables with characteristic function $\phi(t; \theta) = E[e^{itX_1}]$ where $\theta \in \mathbb{R}^k$ is the parameter of the distribution. Let $Z = X_1 + X_2 + \dots + X_n$. Then for which of the following distributions of X_1 would the characteristic function of Z be of the form $\phi(t; \alpha)$ for some $\alpha \in \mathbb{R}^k$?

Options

1. Negative Binomial
2. Geometric
3. Hypergeometric
4. Discrete Uniform

Question Type : **MSQ**

Question ID : **802437597**

Option 1 ID : **8024372385**

Option 2 ID : **8024372386**

Option 3 ID : **8024372387**

Option 4 ID : **8024372388**

Status : **Not Answered**

Chosen Option : --

Q.53 For $n > 1$, let X_1, X_2, \dots, X_n be random variables such that $E(X_i) = 0$ and $E(X_i^2) = 1$ for all i and $E(X_i X_j) = \rho$ for all $i \neq j$. Which of the following statements are true?

Options 1.

1. $\rho = 0$ if and only if X_1, X_2, \dots, X_n are independent
2. $\text{Var}(X_1 + X_2 + \dots + X_n) = n$ if and only if X_1, X_2, \dots, X_n are independent
3. $\text{Var}(X_1 + X_2 + \dots + X_n) = n$ if and only if X_1, X_2, \dots, X_n are pairwise independent
4. $\text{Var}(X_1 + X_2 + \dots + X_n) = n$ if and only if $\rho = 0$

Question Type : **MSQ**

Question ID : **802437594**

Option 1 ID : **8024372373**

Option 2 ID : **8024372374**

Option 3 ID : **8024372375**

Option 4 ID : **8024372376**

Status : **Not Answered**

Chosen Option : --

Q.54

Let the pdf of X be $f(x|\theta) = \frac{2x}{\theta^2}$ for $0 < x < \theta$, where $\theta > 0$ is unknown parameter. Which of the following are $100(1 - \alpha)\%$ confidence intervals for θ ?

Options

1. $\left[X, \frac{X}{\sqrt{\alpha}} \right]$
2. $[X, 2X]$
3. $\left[\frac{\sqrt{2}}{\sqrt{2-\alpha}} X, \frac{\sqrt{2}}{\sqrt{\alpha}} X \right]$
4. $[0, X]$

Question Type : **MSQ**Question ID : **802437600**Option 1 ID : **8024372397**Option 2 ID : **8024372398**Option 3 ID : **8024372399**Option 4 ID : **8024372400**Status : **Not Answered**

Chosen Option : --

Q.55

Let X and Y be independent random variables with $E(X) = E(Y) = 0$ and $Var(X) = Var(Y) = 1$.

Let $Z = X + Y$. Which of the following statements are correct?

Options

1. $P(|Z| > \varepsilon) \leq 2/\varepsilon^2$
2. $E(|Z|) \leq \sqrt{2}$
3. $E(Z^2) = 2$
4. $P(Z \leq 0) = P(Z \geq 0)$

Question Type : **MSQ**Question ID : **802437593**Option 1 ID : **8024372369**Option 2 ID : **8024372370**Option 3 ID : **8024372371**Option 4 ID : **8024372372**Status : **Not Answered**

Chosen Option : --

Q.56 Consider the Gauss-Markov model $Y_{n \times 1} = X_{n \times p} \beta_{p \times 1} + \varepsilon_{n \times 1}$, where $E(\varepsilon) = \mathbf{0}$ and Dispersion $(\varepsilon) = \sigma^2 I_{n \times n}$. Suppose that $p < n$. Which of the following are correct?

Options

1. Least-squares estimate of β is unique
2. Least-squares estimate of an estimable linear function of β is unique
3. Least-squares estimate of $X\beta$ is unique
4. Determinant $(X^T X) > 0$

Question Type : **MSQ**

Question ID : **802437603**

Option 1 ID : **8024372409**

Option 2 ID : **8024372410**

Option 3 ID : **8024372411**

Option 4 ID : **8024372412**

Status : **Not Answered**

Chosen Option : --

Q.57 Let X_1, X_2, X_3 be a random sample from the uniform distribution on the interval $(0, \theta)$. Suppose the prior distribution of θ is uniform on the interval $(0, 1)$. Let $X_{(3)} = \max\{X_1, X_2, X_3\}$. Consider the squared error loss function. Which of the following statements are necessarily true?

Options

1. Bayes estimator of θ is unique
2. $\frac{1}{X_{(3)}}$ is a Bayes estimator of θ
3. $X_{(3)}$ is a Bayes estimator of θ
4. $\frac{1-X_{(3)}}{X_{(3)}}$ is a Bayes estimator of θ

Question Type : **MSQ**

Question ID : **802437602**

Option 1 ID : **8024372405**

Option 2 ID : **8024372406**

Option 3 ID : **8024372407**

Option 4 ID : **8024372408**

Status : **Not Answered**

Chosen Option : --

Q.58

Consider a Markov chain with transition probability matrix

$$P = \begin{pmatrix} 0 & 0 & 1/2 & 0 & 1/2 \\ 0 & 1/2 & 0 & 1/2 & 0 \\ 1/2 & 0 & 0 & 0 & 1/2 \\ 0 & 1/2 & 0 & 1/2 & 0 \\ 1/2 & 0 & 1/2 & 0 & 0 \end{pmatrix}$$

$$\text{Let } \pi = \left(\frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5} \right).$$

Then which of the following statements are correct?

Options

1. π is a stationary distribution
2. If η is a stationary distribution, then $\eta = \pi$
3. The Markov chain is periodic
4. The Markov chain is irreducible

Question Type : MSQ

Question ID : 802437596

Option 1 ID : 8024372381

Option 2 ID : 8024372382

Option 3 ID : 8024372383

Option 4 ID : 8024372384

Status : Not Answered

Chosen Option : --

Q.59

Let X_1, X_2, \dots, X_n be i.i.d. with the common pdf $f(x|\theta) = \frac{\theta}{x^{\theta+1}}$, for $x > 1$ where $\theta > 1$ is an unknown parameter. Which of the following estimators of θ are consistent?

Options

1. $\frac{1}{n} \sum_{i=1}^n X_i$
2. $\frac{1}{n} \sum_{i=1}^n \log(X_i)$
3. $\frac{n}{\sum_{i=1}^n X_i}$
4. $\frac{n}{\sum_{i=1}^n \log(X_i)}$

Question Type : MSQ

Question ID : 802437598

Option 1 ID : 8024372389

Option 2 ID : 8024372390

Option 3 ID : 8024372391

Option 4 ID : 8024372392

Status : Not Answered

Chosen Option : --

Q.60 For circular systematic sampling, which of the following are correct?

Options 1.

Sample mean is an unbiased estimate for population mean but sample variance is not an unbiased estimate for population variance

2.

Sample mean and sample variance are unbiased estimates for population mean and population variance respectively

3.

Sample mean is not an unbiased estimate for population mean but sample variance is an unbiased estimate for population variance

4.

Neither sample mean nor sample variance is an unbiased estimate for their population counterparts

Question Type : **MSQ**

Question ID : **802437606**

Option 1 ID : **8024372421**

Option 2 ID : **8024372422**

Option 3 ID : **8024372423**

Option 4 ID : **8024372424**

Status : **Not Answered**

Chosen Option : **--**

UGC CSIR NET NOVEMBER 2020

Exam Date : 26.11.2020

Final Answer Key on which result compiled on 28.12.2020

Shift : Second

Subject : (704) Mathematical Sciences

Question ID	Correct Option ID	Question ID	Correct Option ID	Question ID	Correct Option ID
802437491	8024371962	802437546	8024372183		8024372298
802437492	8024371965	802437547	8024372188	802437576	8024372301,
802437493	8024371971	802437548	-----		8024372302,
802437494	8024371976	802437549	8024372195		8024372303,
802437495	8024371980	802437550	8024372200		8024372304
802437496	8024371984	802437551	8024372202,	802437577	8024372305,
802437497	8024371986		8024372203		8024372306,
802437498	8024371991	802437552	8024372205,		8024372307
802437499	8024371996		8024372208	802437578	8024372309,
802437500	8024371999	802437553	8024372209,		8024372310
802437501	8024372004		8024372212	802437579	8024372313,
802437502	8024372007	802437554	8024372213,		8024372315
802437503	8024372010		8024372216	802437580	8024372318,
802437504	8024372013	802437555	8024372217,		8024372320
802437505	8024372017		8024372218,	802437581	8024372322,
802437506	8024372022		8024372219,		8024372323
802437507	8024372027		8024372220	802437582	8024372325,
802437508	8024372029,	802437556	8024372222		8024372327
	8024372031	802437557	8024372226,	802437583	8024372330,
802437509	8024372036		8024372227,		8024372331
802437510	8024372039		8024372228	802437584	8024372334,
802437511	8024372043	802437558	8024372230,		8024372335
802437512	8024372047		8024372231,	802437585	8024372338,
802437513	8024372052		8024372232		8024372339
802437514	8024372056	802437559	8024372234	802437586	8024372341,
802437515	8024372060	802437560	8024372237,		8024372343,
802437516	8024372063		8024372238,		8024372344
802437517	8024372067		8024372239	802437587	8024372345,
802437518	8024372071	802437561	8024372242,		8024372347
802437519	8024372074		8024372243	802437588	8024372350
802437520	8024372080	802437562	8024372245,	802437589	8024372354,
802437521	8024372084		8024372246		8024372355
802437522	8024372086	802437563	8024372250,	802437590	8024372358,
802437523	8024372090		8024372252		8024372360
802437524	8024372094	802437564	8024372253,	802437591	8024372361,
802437525	8024372097		8024372255		8024372363
802437526	8024372101	802437565	8024372257,	802437592	8024372365,
802437527	8024372107		8024372258		8024372367
802437528	8024372107	802437566	8024372261,	802437593	8024372369,
802437529	8024372111		8024372262,		8024372370,
802437530	8024372116		8024372263,		8024372371
802437531	8024372119		8024372264	802437594	8024372376
802437532	8024372123	802437567	8024372265,	802437595	8024372378,
802437533	8024372128		8024372266,		8024372379
802437534	8024372131		8024372268	802437596	8024372381
802437535	8024372133	802437568	8024372270,	802437597	8024372385
802437536	8024372139		8024372271	802437598	8024372392
802437537	8024372144	802437569	8024372276	802437599	8024372396
802437538	8024372147	802437570	8024372278	802437600	8024372397,
802437539	8024372152		8024372281,		8024372399
802437540	8024372153	802437571	8024372282	802437601	8024372401,
802437541	8024372159		8024372287		8024372403
802437542	8024372163	802437572	8024372290,	802437602	8024372405,
802437543	8024372166		8024372291		8024372408
802437544	8024372171	802437573	8024372293,	802437603	8024372410,
802437545	8024372173		8024372295		8024372411
	8024372177	802437574	8024372297,	802437604	8024372413,
		802437575			

NB:-Correction option ID ----- means the question remains cancelled and marks will be given to only those candidates who attempted that question Page 1

Question ID	Correct Option ID
-------------	-------------------

	8024372414,
	8024372415
802437605	8024372417,
	8024372418,
	8024372419,
	8024372420
802437606	8024372421
802437607	8024372425,
	8024372426,
	8024372427
802437608	8024372430,
	8024372431
802437609	8024372433,
	8024372434,
	8024372435,
	8024372436
802437610	8024372438,
	8024372439,
	8024372440

P Kalika Maths

Some Useful Links:

- 1. Free Maths Study Materials** (<https://pkalika.in/2020/04/06/free-maths-study-materials/>)
- 2. BSc/MSc Free Study Materials** (<https://pkalika.in/2019/10/14/study-material/>)
- 3. MSc Entrance Exam Que. Paper:** (<https://pkalika.in/2020/04/03/msc-entrance-exam-paper/>)
[JAM(MA), JAM(MS), BHU, CUCET, ...etc]
- 4. PhD Entrance Exam Que. Paper:** (<https://pkalika.in/que-papers-collection/>)
[CSIR-NET, GATE(MA), BHU, CUCET,IIT, NBHM, ...etc]
- 5. CSIR-NET Maths Que. Paper:** (<https://pkalika.in/2020/03/30/csir-net-previous-yr-papers/>)
[Upto 2019 Dec]
- 6. Practice Que. Paper:** (<https://pkalika.in/2019/02/10/practice-set-for-net-gate-set-jam/>)
[Topic-wise/Subject-wise]
- 7. List of Maths Suggested Books** (<https://pkalika.in/suggested-books-for-mathematics/>)
- 8. CSIR-NET Mathematics Details Syllabus** (<https://wp.me/p6gYUB-Fc>)
- 9. Free Video Lectures for CSIR-NET, GATE, SET, Asst. Prof. ..etc**
<https://www.youtube.com/c/PKALIKA>